




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 217/04, 401/12, A61K 31/47		A1	(11) International Publication Number: WO 98/51671
			(43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/EP98/02584		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 28 April 1998 (28.04.98)			
(30) Priority Data: 9709303.3 9 May 1997 (09.05.97) GB		Published With international search report.	
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): JOHNSON, Christopher, Norbert [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). STEMPE, Geoffrey [GB/GB]; SmithKline Beecham Pharmaceuticals, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).			
(74) Agent: GARRETT, Michael; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).			

(54) Title: SUBSTITUTED TETRAHYDROISOQUINOLINE DERIVATIVES AS MODULATORS OF DOPAMINE D3 RECEPTORS



(57) Abstract

Compounds of formula (I) wherein: R¹ represents a substituent selected from: a hydrogen or halogen atom; a hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, trifluoromethanesulfonyloxy, pentafluoroethyl, C₁₋₄alkyl, C₁₋₄alkoxy, arylC₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkoxy C₁₋₄alkyl, C₃₋₆cycloalkylC₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulphonyl, C₁₋₄alkylsulphonyloxy, C₁₋₄alkylsulphonylC₁₋₄alkyl, arylsulphonyl arylsulphonyloxy, arylsulphonylC₁₋₄alkyl, C₁₋₄alkylsulphonamido, C₁₋₄alkylamido, C₁₋₄alkylsulphonamidoC₁₋₄alkyl, C₁₋₄alkylamidoC₁₋₄alkyl, arylsulphomanido, arylcarboxamido, arylsulphonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aryl, arylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group; a group R³OCO(CH₂)_p, R³CON(R⁴)(CH₂)_p, R³R⁴NCO(CH₂)_p, or R³R⁴NSO(CH₂)_p where each of R³ and R⁴ independently represents a hydrogen atom or a C₁₋₄alkyl group or R³R⁴ forms part of a C₃₋₆azacycloalkane or C₃₋₆(2-oxo)azacycloalkane ring and p represents zero or an integer from 1 to 4; or a group Ar³-Z, wherein Ar³ represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring and Z represents a bond O, S, or CH₂; s represents an integer from zero to 2 and r represents an integer from 1 to 4, such that the sum of s + r is 1 to 4; t represents an integer from zero to 1 and u represents an integer from zero to 2; R² represents a hydrogen atom or a C₁₋₄alkyl group; q is 1 or 2; A represents a group of the formula (a), (b) or (c): wherein Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; or an optionally substituted bicyclic ring system; Ar¹ and Ar² each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; and Y represents a bond, -NHCO-, -CONH-, -CH₂-, or -(CH₂)_nY'(CH₂)_m-, wherein Y' represents O, S, SO₂, or CO and m and n each represents zero or 1 such that the sum of m+n is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar ortho to the carboxamide moiety is necessarily a hydrogen or methoxy group; and salts thereof. Compounds of formula (I) and their salts have affinity for dopamine receptors, in particular the D₃ receptor, and thus potential in the treatment of conditions wherein modulation of the D₃ receptor is beneficial, e.g. as antipsychotic agents.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						


SUBSTITUTED TETRAHYDROISOQUINOLINE DERIVATIVES AS MODULATORS OF DOPAMINE D₃ RECEPTORS

The present invention relates to novel tetrahydroisoquinoline derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy, as modulators of dopamine D₃ receptors, in particular as antipsychotic agents.


US Patent No. 5,294,621 describes tetrahydropyridine derivatives of the formula:



10

wherein  is an optionally substituted thienyl or optionally substituted phenyl ring; R¹, R² and R³ are each *inter alia* hydrogen; X is *inter alia* (CH₂)_mNR⁷CO; m is 2-4; and Ar¹ is an optionally substituted heterocyclic ring or an optionally substituted phenyl ring. The compounds are said to be useful as antiarrhythmic agents.

15 European Patent Application 0 464 846 A1 describes imide derivatives of the formula:




wherein B is a carbonyl group or a sulphonyl group, R¹, R², R³ and R⁴ are each hydrogen or a lower alkyl group, or R¹ and R² or R¹ and R³ may be combined together to make a non-aromatic hydrocarbon ring, or R¹ and R³ may be combined together to make an aromatic ring, and n is 0 or 1; A is a non-aromatic hydrocarbon ring, and p and q are each 0, 1, or 2; Ar is an aromatic ring, a heteroaromatic group, a benzoyl group, a phenoxy group, or a phenylthio group and G is N, CH, or CHOH. The compounds are said to be useful as antipsychotic agents.

25 WO 95/10513 describes benzothiophene derivatives and related compounds as estrogen agonists.

We have now found a class of tetrahydroisoquinoline derivatives which have affinity for dopamine receptors, in particular the D₃ receptor, and thus potential in the treatment of conditions wherein modulation of the D₃ receptor is beneficial, eg as antipsychotic agents.

30 In a first aspect the present invention provides compounds of formula (I) :



Formula (I)

wherein:

R¹ represents a substituent selected from: a hydrogen or halogen atom; a hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, trifluoromethanesulfonyloxy, pentafluoroethyl, C₁₋₄alkyl, C₁₋₄alkoxy, arylC₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkoxyC₁₋₄alkyl, C₃₋₆cycloalkylC₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulphonyl, C₁₋₄alkylsulphonyloxy, C₁₋₄alkylsulphonylC₁₋₄alkyl, arylsulphonyl, arylsulphonyloxy, arylsulphonylC₁₋₄alkyl, C₁₋₄alkylsulphonamido, C₁₋₄alkylamido, C₁₋₄alkylsulphonamidoC₁₋₄alkyl, C₁₋₄alkylamidoC₁₋₄alkyl, arylsulphonamido, arylcarboxamido, arylsulphonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aroyl, aroylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group; a group R³OCO(CH₂)_p, R³CON(R⁴)(CH₂)_p, R³R⁴NCO(CH₂)_p or R³R⁴NSO₂(CH₂)_p where each of R³ and R⁴ independently represents a hydrogen atom or a C₁₋₄alkyl group or R³R⁴ forms part of a C₃₋₆azacycloalkane or C₃₋₆(2-oxo)azacycloalkane ring and p represents zero or an integer from 1 to 4; or a group Ar³-Z, wherein Ar³ represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring and Z represents a bond, O, S, or CH₂;



R² represents a hydrogen atom or a C₁₋₄alkyl group;

q is 1 or 2;

s represents an integer from zero to 2 and r represents an integer from 1 to 4, such that the sum of s + r is 1 to 4;

t represents an integer from zero to 1 and u represents an integer from zero to 2;

A represents a group of the formula (a), (b) or (c):



wherein

Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; or an optionally substituted bicyclic system;

Ar¹ and Ar² each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; and

Y represents a bond, -NHCO-, -CONH-, -CH₂-, or -(CH₂)_mY¹(CH₂)_n-, wherein Y¹ represents O, S, SO₂, or CO and m and n each represent zero or 1 such that the sum of m+n is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar *ortho* to the carboxamide moiety is necessarily a hydrogen or a methoxy group;

and salts thereof.

In the compounds of formula (I) above an alkyl group or moiety may be straight or branched. Alkyl groups which may be employed include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl and any branched isomers thereof such as isopropyl, t-butyl, sec-pentyl, and the like.

Examples of compounds of formula (I) include those in which Ar is a bicyclic aromatic or heteroaromatic ring system, and t and u are both 1 and in which R¹ is other than pentafluoroethyl.

When R¹ represents an arylC₁₋₄alkoxy, arylsulphonyl, arylsulphonyloxy, arylsulphonylC₁₋₄alkyl, arylsulphonamido, arylcarboxamido, arylsulphonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aroyl, aroylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group, the aryl moiety may be selected from an optionally substituted phenyl ring or an optionally substituted 5- or 6-membered heterocyclic ring. In the group R¹ an aryl moiety may be optionally substituted by one or more substituents selected from hydrogen, halogen, amino, cyano, C₁₋₄alkyl, C₁₋₄alkylamino, C₁₋₄dialkylamino, C₁₋₄alkylamido, C₁₋₄alkanoyl, or R⁵R⁶NCO where each of R⁵ and R⁶ independently represents a hydrogen atom or C₁₋₄alkyl group.

A halogen atom present in the compounds of formula (I) may be fluorine, chlorine, bromine or iodine.

When q is 2, the substituents R¹ may be the same or different.

An optionally substituted 5- or 6-membered heterocyclic aromatic ring, as defined for any of the groups Ar, Ar¹, Ar² or Ar³ may contain from 1 to 4 heteroatoms selected from O, N or S. When the ring contains 2-4 heteroatoms, one is preferably selected from O, N and S and the remaining heteroatoms are preferably N. Examples of 5 and 6-membered heterocyclic groups include furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, pyridyl, triazolyl, triazinyl, pyridazyl, pyrimidinyl and pyrazolyl.

Examples of bicyclic, for example, bicyclic aromatic or heteroaromatic, ring systems for Ar include naphthyl, indazolyl, indolyl, benzofuranyl, benzothienyl, benzothiazolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzisothiazolyl, quinolinyl, quinoxolinyl, quinazolinyl, cinnolinyl, isoquinolinyl, pyrazolo[1,5-a]pyrimidyl, pyrrolo[3,2-b]pyridyl, pyrrolo[3,2-c]pyridyl, thieno[3,2-b]thiophenyl, 1,2-dihydro-2-oxo-quinolinyl, 2,3-dihydro-3-oxo-4H-benzoxazinyl, 1,2-dihydro-2-oxo-3H-indolyl.

The rings Ar, Ar¹, or Ar² may each independently be optionally substituted by one or more substituents selected from: a hydrogen or halogen atom, or a hydroxy, cyano, nitro, C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylenedioxy, C₁₋₄alkanoyl, C₁₋₄alkylsulphonyl, C₁₋₄alkylsulphinyl, C₁₋₄alkylthio, R⁷SO₂N(R⁸)-, R⁷R⁸N-, R⁷R⁸NCO-, R⁷R⁸NSO₂-, or R⁷CON(R⁸)- group wherein each of R⁷ and R⁸ independently represents a hydrogen atom or a C₁₋₄ alkyl group, or R⁷R⁸ together form a C₃₋₆ alkylene chain.

Alternatively, Ar and Ar² may be optionally substituted by one or more 5- or 6- membered heterocyclic rings, as defined above, optionally substituted by a C₁₋₂ alkyl or R⁷R⁸N- group; wherein R⁷ and R⁸ are as defined above.

5 In the rings Ar and Ar² substituents positioned *ortho* to one another may be linked to form a 5- or 6- membered ring.

It will be appreciated that for use in medicine the salts of formula (I) should be physiologically acceptable. Suitable physiologically acceptable salts will be apparent to those skilled in the art and include for example acid addition salts formed with inorganic acids eg. hydrochloric, hydrobromic, sulphuric, nitric or phosphoric acid; and organic
10 acids eg. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulphonic, methanesulphonic or naphthalenesulphonic acid. Other non-physiologically acceptable salts eg. oxalates, may be used, for example in the isolation of compounds of formula (I) and are included within the scope of this invention. Also included within the scope of the invention are solvates and hydrates of compounds of formula (I).

15 Certain of the compounds of formula (I) may form acid addition salts with one or more equivalents of the acid. The present invention includes within its scope all possible stoichiometric and non-stoichiometric forms.

It will be appreciated certain of the compounds of formula (I) contain two asymmetric centres. Such compounds can exist in diastereomeric forms, namely *cis*- and
20 *trans*- isomers; both forms and all mixtures thereof are included within the scope of this invention. Furthermore, each diastereoisomer can exist as optical isomers (enantiomers). Both the pure enantiomers, racemic mixtures (50% of each enantiomer) and unequal mixtures of the two are included within the scope of the invention. In accordance with convention the (+) and (-) designations used herein indicate the direction of rotation of
25 plane-polarised light by the compounds. The prefix (+) indicates that the isomer is dextrorotatory (which can also be designated d) and the prefix (-) indicates the levorotatory isomer (which can also be designated l). It will thus be appreciated that the invention extends to the individual diastereoisomers, individual enantiomers and any and all mixtures of these forms.

30 Certain of the other compounds of formula (I) can also exist in the form of *cis*- and *trans*- isomers. The present invention includes within its scope all such isomers, including mixtures.


In compounds of formula (I), it is preferred that either t and u are both zero or that t and u are both 1.

35 Certain of the substituted heteroaromatic ring systems included in compounds of formula (I) may exist in one or more tautomeric forms. The present invention includes within its scope all such tautomeric forms, including mixtures.

Particular compounds according to the invention include those specifically exemplified and named hereinafter.

40 The present invention also provides a process for preparing compounds of formula (I) which process comprises:

(a) reacting a compound of formula (V):



Formula (V)

5 with a compound of formula (VI):

A-COX


Formula (VI)

10

wherein A is as hereinbefore defined and X is a halogen atom or the residue of an activated ester;

(b) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is a bond, reacting a compound of formula (VII):

15




Formula (VII)

20 wherein one R¹ᵃ represents a group W wherein W is a halogen atom or a trifluoromethylsulphonyloxy group, or W is a group M selected from a boron derivative e.g. a boronic acid function B(OH)₂ or a metal function such as trialkylstannyl e.g. SnBu₃, zinc halide or magnesium halide, and when q is 2 the other R¹ᵃ is R¹; with a compound Ar³-W¹, wherein W¹ is a halogen atom or a trifluoromethylsulphonyloxy group when W is a group M or W¹ is a group M when W is a halogen atom or a trifluoromethylsulphonyloxy group;

25

(c) to prepare a compound of formula (I) wherein R¹ is Ar³-Z and Z is O or S, reacting a compound of formula (VIII):




30

Formula (VIII)

wherein one R¹ᵇ represents a group ZH and when q is 2 the other R¹ᵇ represents R¹; with a reagent serving to introduce the group Ar³;

(d) to prepare a compound of formula (I) where Y is a bond, reaction of a compound of formula (IX):



Formula (IX)


wherein R^1 , R^2 , Ar^1 and W are as hereinbefore defined, with a compound Ar^2-W^1 , wherein W^1 is a halogen atom or a trifluoromethylsulphonyloxy group when W is a group M , or W^1 is a group M when W is a halogen atom or a trifluoromethylsulphonyloxy group.

(e) interconversion of one compound of formula (I) to a different compound of formula (I) e.g. (i) alkylation of a compound (I) wherein R^2 represents hydrogen, (ii) conversion of one R^1 from alkoxy (e.g. methoxy) to hydroxy, or (iii) conversion of R^1 from hydroxy to sulphonyloxy, eg alkylsulphonyloxy or trifluoromethanesulphonyloxy; (iv) conversion of a compound wherein Y represents S to a compound wherein Y is SO_2 or (v) conversion of Y from CO to CH_2 ;

(f) where appropriate, separation of enantiomers, diastereoisomers, or *cis*- and *trans*- isomers of compounds of formula (I), or intermediates thereto, by conventional methods, e.g. chromatography or crystallisation; and optionally thereafter forming a salt of formula (I).

Compounds of formula (V) may be prepared by :-

(g) conversion of a compound of formula (IV):




Formula (IV)


wherein R^1 , R^2 , r , s , t and u are as hereinbefore defined and P is a protecting group such as *t*-butoxycarbonyl or trifluoroacetyl, to a compound of formula (V).

Compounds of formula (IV) in which t is 1 may be prepared by:-

(h) by reacting a compound of formula (II):

**Formula (II)**

- 5 wherein R¹ and q are as hereinbefore defined; with a compound of formula (IIIa):


**Formula (IIIa)**

10

wherein P, R², r, s, and u are as hereinbefore defined;

Compounds of formula (IV) where t is zero may be prepared by: -

- 15 (i) reacting a compound of formula (II), wherein R¹ and q are hereinbefore defined, with a compound of formula (IIIb):


**Formula (IIIb)**

20

wherein P, R², r, s, and u are as hereinbefore defined.

Compounds of formula (V), where t and u are both zero may be prepared by:-

- 25 (j) conversion of a compound of formula (X):-


**Formula (X)**

- 30 wherein R¹, q, r and s are as hereinbefore defined and v is 1 or 2, into a corresponding ketone, followed by reductive amination. This may be effected by methods well known

in the art for (i) conversion of a ketal to a ketone in the presence of aqueous acid; followed by (ii) reductive amination of the ketone with R^2NH_2 or ammonium acetate in the presence of a reducing agent. Suitable reducing agents which may be employed include sodium borohydride, cyanoborohydride or triacetoxyborohydride under acidic conditions, or catalytic hydrogenation. The reaction may conveniently be effected in a solvent such as methanol, ethanol or dichloroethane..

Compounds of formula (X) wherein R^1 and q are as hereinbefore defined, may be prepared by:-

(k) reacting a compound of formula (XI):-



Formula (XI)

wherein v, r and s are as hereinbefore defined, with a compound of formula (II), wherein R^1 and q are as hereinbefore defined.

Processes (h), (i) and (k) require the presence of a reducing agent. Suitable reducing agents which may be employed include sodium borohydride, cyanoborohydride or triacetoxyborohydride under acidic conditions, or catalytic hydrogenation. The reaction may conveniently be effected in a solvent such as ethanol.

Process (g) may be effected by standard methods well known in the art for (i) removal of a t-butoxycarbonyl group, e.g., using acidic conditions; (ii) removal of a trifluoroacetyl group, e.g., using basic conditions.

Reaction of a compound of formula (VII) with Ar^3W^1 , according to process (b) or a compound of formula (IX) with Ar^2-W^1 according to process (d) may be effected in the presence of a transition metal eg palladium catalyst such as *bis*-triphenylphosphinepalladium dichloride or *tetrakis*-triphenylphosphinepalladium (0). When M represents a boronic acid function such as $B(OH)_2$, the reaction may be carried out under basic conditions, for example using aqueous sodium carbonate in a suitable solvent such as dioxane. When M is trialkylstannyl the reaction may be carried out in an inert solvent, such as xylene or dioxane optionally in the presence of LiCl. When M is a zinc or magnesium halide the reaction may be effected in an aprotic solvent such as tetrahydrofuran. The substituent W is preferably a halogen atom such as bromine, or a sulphonyloxy group such as trifluoromethylsulphonyloxy; and W^1 is preferably a group M, such as trialkylstannyl or $B(OH)_2$.

In process (c) the reagent serving to introduce the group Ar^3 is preferably a compound of formula Ar^3-Hal , wherein Hal is a halogen atom. The reaction may be effected in the presence of a base, such as potassium carbonate, in a solvent such as dimethylformamide.

Interconversion reactions according to process (e) may be effected using methods well known in the art.

Compounds of formula (II) may be prepared by methods known in the art.

Compounds of formula (IIIa) and (IIIb) are known or may be prepared using
5 standard procedures.

Compounds of formula (VII), (VIII) or (IX) may be prepared by processes analogous to (a), (g), (h) and (i) described above. Compounds Ar^2W^1 , Ar^3W^1 and Ar^3Hal are commercially available or may be prepared by standard methods.

Compounds of formula (XI) are commercially available or may be prepared using
10 standard procedures.

Compounds of formula (I) have been found to exhibit affinity for dopamine receptors, in particular the D_3 receptor, and are expected to be useful in the treatment of disease states which require modulation of such receptors, such as psychotic conditions. Compounds of formula (I) have also been found to have greater affinity for dopamine D_3
15 than for D_2 receptors. The therapeutic effect of currently available antipsychotic agents (neuroleptics) is generally believed to be exerted via blockade of D_2 receptors; however this mechanism is also thought to be responsible for undesirable extrapyramidal side effects (eps) associated with many neuroleptic agents. Without wishing to be bound by theory, it has been suggested that blockade of the recently characterised dopamine D_3
20 receptor may give rise to beneficial antipsychotic activity without significant eps. (see for example Sokoloff et al, Nature, 1990; 347: 146-151; and Schwartz et al, Clinical Neuropharmacology, Vol 16, No. 4, 295-314, 1993). Preferred compounds of the present invention are therefore those which have higher affinity for dopamine D_3 than dopamine D_2 receptors (such affinity can be measured using standard methodology for
25 example using cloned dopamine receptors). Said compounds may advantageously be used as selective modulators of D_3 receptors.

We have found that certain compounds of formula (I) are dopamine D_3 receptor antagonists, others may be agonists or partial agonists. The functional activity of compounds of the invention (i.e. whether they are antagonists, agonists or partial
30 agonists) can be readily determined using the test method described hereinafter, which does not require undue experimentation. D_3 antagonists are of potential use as antipsychotic agents for example in the treatment of schizophrenia, schizo-affective disorders, psychotic depression, mania, paranoid and delusional disorders. Conditions which may be treated by dopamine D_3 receptor agonists include dyskinetic disorders
35 such as Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias; depression; anxiety, memory disorders, sexual dysfunction and drug (eg. cocaine) dependency.

In a further aspect therefore the present invention provides a method of treating conditions which require modulation of dopamine D_3 receptors, for example psychoses
40 such as schizophrenia, which comprises administering to a subject in need thereof an effective amount of a compound of formula (I) or a physiologically acceptable salt thereof.

The invention also provides the use of a compound of formula (I) or a physiologically acceptable salt thereof in the manufacture of a medicament for the treatment of conditions which require modulation of dopamine D₃ receptors, for example psychoses such as schizophrenia.

5 A preferred use for D₃ antagonists according to the present invention is in the treatment of psychoses such as schizophrenia.

A preferred use for D₃ agonists according to the present invention is in the treatment of dyskinetic disorders such as Parkinson's disease.

10 For use in medicine, the compounds of the present invention are usually administered as a standard pharmaceutical composition. The present invention therefore provides in a further aspect pharmaceutical compositions comprising a novel compound of formula (I) or a physiologically acceptable salt thereof and a physiologically acceptable carrier.

15 The compounds of formula (I) may be administered by any convenient method, for example by oral, parenteral, buccal, sublingual, nasal, rectal or transdermal administration and the pharmaceutical compositions adapted accordingly.

The compounds of formula (I) and their physiologically acceptable salts which are active when given orally can be formulated as liquids or solids, for example syrups, suspensions or emulsions, tablets, capsules and lozenges.

20 A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable liquid carrier(s) for example an aqueous solvent such as water, ethanol or glycerine, or a non-aqueous solvent, such as polyethylene glycol or an oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

25 A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and cellulose.

30 A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatin capsule; alternatively, a dispersion or suspension can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatin capsule.

35 Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

40 Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an

atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a propellant which can be a compressed gas such as compressed air or an organic propellant such as a fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base such as cocoa butter.

Compositions suitable for transdermal administration include ointments, gels and patches.

Preferably the composition is in unit dose form such as a tablet, capsule or ampoule.

Each dosage unit for oral administration contains preferably from 1 to 250 mg (and for parenteral administration contains preferably from 0.1 to 25 mg) of a compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base.

The physiologically acceptable compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1 mg and 500 mg, preferably between 10 mg and 400 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

Biological Test Methods

The ability of the compounds to bind selectively to human D₃ dopamine receptors can be demonstrated by measuring their binding to cloned receptors. The inhibition constants (K_i) of test compounds for displacement of [¹²⁵I] iodosulpride binding to human D₃ dopamine receptors expressed in CHO cells were determined as follows. The cell lines were shown to be free from bacterial, fungal and mycoplasmal contaminants, and stocks of each were stored frozen in liquid nitrogen. Cultures were grown as monolayers or in suspension in standard cell culture media. Cells were recovered by scraping (from monolayers) or by centrifugation (from suspension cultures), and were washed two or three times by suspension in phosphate buffered saline followed by collection by centrifugation. Cell pellets were stored frozen at -40°C. Crude cell membranes were prepared by homogenisation followed by high-speed centrifugation, and characterisation of cloned receptors achieved by radioligand binding.

Preparation of CHO cell membranes

Cell pellets were gently thawed at room temperature, and resuspended in about 20 volumes of ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), 20mM EDTA, 0.2 M sucrose.

- 5 The suspension was homogenised using an Ultra-Turrax at full speed for 15 sec. The homogenate was centrifuged at 18,000 r.p.m for 20 min at 4°C in a Sorvall RC5C centrifuge. The membrane pellet was resuspended in ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), using an Ultra-Turrax, and recentrifuged at 18,000 r.p.m for 15 min at 4°C in a Sorvall RC5C. The membranes were washed two more times with ice-cold 50 mM Tris salts (pH 7.4 @ 37°C). The final pellet was resuspended in 50 mM Tris salts (pH 7.4 @ 37°C), and the protein content determined using bovine serum albumin as a standard (Bradford, M. M. (1976) Anal. Biochem. 72, 248-254).

Binding experiments on cloned dopamine receptors

- 15 Crude cell membranes were incubated with 0.1 nM [¹²⁵I] iodosulpride (~2000 Ci/mmol; Amersham, U. K.), and the test compound in a buffer containing 50 mM Tris salts (pH 7.4 @ 37°C), 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂, 0.1% (w/v) bovine serum albumin, in a total volume of 1 ml for 30 min at 37°C. Following incubation, samples were filtered using a Brandel Cell Harvester, and washed three times with ice-cold 50 mM Tris salts (pH 7.4 @ 37°C), 120 mM NaCl, 5 mM KCl, 2 mM CaCl₂, 1 mM MgCl₂. The radioactivity on the filters was measured using a Cobra gamma counter (Canberra Packard). Non-specific binding was defined as the radioligand binding remaining after incubation in the presence of 100 µM iodosulpride. For competition curves, 14 concentrations (half-log dilutions) of competing cold drug were used.
- 20 Competition curves were analysed simultaneously whenever possible using non-linear least-squares fitting procedures, capable of fitting one, two or three site models.

Compounds of Examples tested according to this method had pK_i values in the range 7.0 - 8.5 at the human cloned dopamine D₃ receptor.

Functional Activity at cloned dopamine receptors

- The functional activity of compounds at human D₂ and human D₃ receptors (ie agonism or antagonism) may be determined using a Cytosensor Microphysiometer (McConnell HM et al Science 1992 257 1906-1912) In Microphysiometer experiments, cells
- 35 (hD₂_CHO or hD₃_CHO) were seeded into 12mm Transwell inserts (Costar) at 300000 cells/cup in foetal calf serum (FCS)-containing medium. The cells were incubated for 6h at 37°C in 5%CO₂, before changing to FCS-free medium. After a further 16-18h, cups were loaded into the sensor chambers of the Cytosensor Microphysiometer (Molecular Devices) and the chambers perfused with running medium (bicarbonate-free Dulbecco's modified Eagles medium containing 2 mM glutamine and 44 mM NaCl) at a flow rate of
- 40 100 µl/min. Each pump cycle lasted 90s. The pump was on for the first 60s and the acidification rate determined between 68 and 88s, using the Cytosoft programme. Test compounds were diluted in running medium. In experiments to determine agonist

activity, cells were exposed (4.5 min for hD2, 7.5 min for hD3) to increasing concentrations of putative agonist at half hour intervals. Seven concentrations of the putative agonist were used. Peak acidification rate to each putative agonist concentration was determined and concentration-response curves fitted using Robofit [Tilford, N.S.,
 5 Bowen, W.P. & Baxter, G.S. Br. J. Pharmacol. (1995) in press]. In experiments to determine antagonist potency, cells were treated at 30 min intervals with five pulses of a submaximal concentration of quinpirole (100 nM for hD2 cells, 30 nM for hD3 cells), before exposure to the lowest concentration of putative antagonist. At the end of the next
 10 30 min interval, cells were pulsed again with quinpirole (in the continued presence of the antagonist) before exposure to the next highest antagonist concentration. In all, five concentrations of antagonist were used in each experiment. Peak acidification rate to each agonist concentration was determined and concentration-inhibition curves fitted using Robofit.

15 **Pharmaceutical Formulations**

The following represent typical pharmaceutical formulations according to the present invention, which may be prepared using standard methods.

IV Infusion

20	Compound of formula (I)	1-40 mg
	Buffer	to pH ca 7
	Solvent/complexing agent	to 100 ml

Bolus Injection

	Compound of formula (I)	1-40 mg
25	Buffer	to pH ca 7
	Co-Solvent	to 5 ml

Buffer : Suitable buffers include citrate, phosphate, sodium hydroxide/hydrochloric acid.

30 Solvent : Typically water but may also include cyclodextrins (1-100 mg) and co-solvents such as propylene glycol, polyethylene glycol and alcohol.

Tablet

	Compound	1 - 40 mg
35	Diluent/Filler *	50 - 250 mg
	Binder	5 - 25 mg
	Disintegrant *	5 - 50 mg
	Lubricant	1 - 5 mg
	Cyclodextrin	1 - 100 mg

40

* may also include cyclodextrins

Diluent : e.g. Microcrystalline cellulose, lactose, starch
 Binder : e.g. Polyvinylpyrrolidone, hydroxypropymethylcellulose
 Disintegrant : e.g. Sodium starch glycollate, crospovidone
 Lubricant : e.g. Magnesium stearate, sodium stearyl fumarate.

5

Oral Suspension

Compound	1 - 40 mg
Suspending Agent	0.1 - 10 mg
Diluent	20 - 60 mg
10 Preservative	0.01 - 1.0 mg
Buffer	to pH ca 5 - 8
Co-solvent	0 - 40 mg
Flavour	0.01 - 1.0 mg
Colourant	0.001 - 0.1 mg

15 Suspending agent : e.g. Xanthan gum, microcrystalline cellulose
 Diluent : e.g. sorbitol solution, typically water
 Preservative : e.g. sodium benzoate
 Buffer : e.g. citrate
 20 Co-solvent : e.g. alcohol, propylene glycol, polyethylene glycol, cyclodextrin

The invention is further illustrated by the following non-limiting examples :

Description 1

25

7-Bromo-1,2,3,4-tetrahydroisoquinoline

A mixture of 7-bromo-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline (G.E. Stokker, Tetrahedron Letters 1996, 37, 5453) (43.4g, 0.14 mol), potassium carbonate (104.3g, 0.75 mol), methanol (1L) and water (150ml) was heated at
 30 reflux for 1h, then cooled and evaporated *in vacuo*. Residue was partitioned between water (1L) and dichloromethane (4 x 200ml). Combined extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give an oil which was dissolved in hexane. The mixture was filtered and the filtrate evaporated *in vacuo* to give the
 35 title compound as an oil (17.7g, 60%).

¹H NMR (CDCl₃) δ: 1.77 (1H, br s), 2.73 (2H, t, J = 7 Hz), 3.13 (2H, t, J = 7 Hz), 3.98 (2H, s), 6.96 (1H, d, J = 9 Hz), 7.16 (1H, d, J = 2 Hz), 7.26 (1H, dd, J = 9, 2 Hz).

40

The following compounds were prepared in a similar manner to Description 1

(a) 7-Cyano-1,2,3,4-tetrahydroisoquinoline

5

Mass spectrum (AP⁺): Found 159 (MH⁺). C₁₀H₁₀N₂ requires 158.

Description 2

10 7-Cyano-2-trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline

A mixture of 7-bromo-2- trifluoroacetyl -1,2,3,4-tetrahydroisoquinoline (51.7 g, 0.168 mol), copper (I) cyanide (31.8 g, 0.35 mol) and N-methyl-2-pyrrolidinone (620 ml) was heated at reflux for 4h, cooled, then partitioned between dilute aqueous ammonia (1.5 L) and dichloromethane (5 x 300ml). The combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound (42.6 g, 100 %) as an oil.

20

Mass spectrum (AP⁺): Found 253 (M-H)⁺. C₁₂H₉F₃N₂O requires 254.

Description 3

(±)-trans-2-((N-(tert-Butyloxycarbonyl)amino)methyl)cyclopropane-1-carboxaldehyde

25 To a solution of (±)-trans-1-((N-(tert-butyloxycarbonyl)amino)methyl)-2-((tert-butyl)diphenylsilyloxy)methyl)cyclopropane [T. Morikawa et al, J. Org. Chem., 1994, 59, 97] (0.33g, 0.75 mmol) in dry THF (10ml) at 0°C, was added a 1M solution of tetra-n-butylammonium fluoride in THF (2.3ml, 2.3 mmol). The mixture was stirred at room temperature for 3 hours, then partitioned between diethyl ether (25ml) and water (25ml).
30 Aqueous phase was further extracted with diethyl ether (25ml x 2) and the combined organic extracts were washed with brine (40ml), dried (Na₂SO₄) then evaporated *in vacuo* to give an oil. To a solution of oxalyl chloride (0.08g, 0.6 mmol) in dry dichloromethane (3ml) at -80°C under argon, was added dropwise a solution of dimethyl sulfoxide (0.09g, 1.2 mmol) in dichloromethane (0.5ml). The resulting mixture was stirred at -78°C for
35 0.75h, then a solution of the above oil in dry dichloromethane (3ml) was added. The mixture was stirred for 1h then triethylamine (1ml) was added and the mixture warmed to room temperature. The mixture was partitioned between dichloromethane (100ml) and water (50ml). The organic layer was washed with water (30ml) and brine (30ml), then dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound as an oil (0.12g, 98%)

40

¹H NMR (CDCl₃) δ: 1.07 (1H, m), 1.30 (1H, m), 1.45 (9H, s), 1.69 - 1.90 (2H, m), 2.95 - 3.30 (2H, m), 4.75 (1H, br s), 9.09 (1H, d, J = 5 Hz).

Description 4

- 5 (±)-*trans*-1-(*N*-(*tert*-Butyloxycarbonyl)amino)methyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)methylcyclopropane

A mixture of (±)-*trans*-2-((*N*-(*tert*-butyloxycarbonyl)amino)cyclopropane-1-carboxaldehyde (0.12g, 0.6 mmol), 7-cyano-1,2,3,4-tetrahydroisoquinoline (0.11g, 0.66
10 mmol) and sodium triacetoxyborohydride (0.19g, 0.9 mmol) in 1,2-dichloromethane (15ml) was allowed to stir at room temperature for 20h, then partitioned between dichloromethane (120ml) and saturated aqueous NaHCO₃ (40ml). Organic phase was washed with saturated NaHCO₃ (40ml), brine (40ml), dried (Na₂SO₄) and evaporated *in vacuo* to an oil. Chromatography on silica with ethylacetate-hexane 20 - 40% gradient
15 elution gave the title compound as an amber oil (0.16g, 78%).

Mass spectrum (API⁺): Found 342 (MH⁺). C₂₀H₂₇N₃O₂ requires 341.

¹H NMR (CDCl₃) δ: 0.40 - 0.55 (2H, m), 0.85 - 0.91 (2H, m), 1.47 (9H, s), 2.34 - 2.60
20 (2H, m), 2.75 - 2.85 (2H, m), 2.90 - 3.00 (2H, m), 3.02 - 3.10 (2H, m), 3.55 - 3.80 (2H, m), 4.68 (1H, br s), 7.20 (1H, d, J = 8 Hz), 7.35 (1H, s), 7.40 (1H, d, J = 8 Hz).

The following compound was prepared in a similar manner to Description 4.

- 25 (a) *trans*-2-(1-(4-(*t*-Butyloxycarbonyl)aminomethyl)cyclohexylmethyl)-7-cyano-1,2,3,4-tetrahydroisoquinoline

Mass spectrum (API⁺): Found 384 (MH⁺). C₂₃H₃₃N₃O₂ requires 383.

30

Description 5

- 35 (±)-*trans*-1-Aminomethyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)-methylcyclopropane

To a solution of (±)-*trans*-1-(*N*-(*tert*-butyloxycarbonyl)methyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)methylcyclopropane (0.16g, 0.47 mmol) in dry dichloromethane (10ml) at 0°C, was added trifluoroacetic acid (0.36ml). The mixture was stirred at 0°C for 1h, then more trifluoroacetic acid (0.4ml) was added. The mixture was stirred at room

temperature for 5h, then partitioned between dichloromethane (100ml) and saturated aqueous NaHCO_3 (50ml). Organic phase was washed with brine (50ml), dried (Na_2SO_4) and evaporated *in vacuo* to give the title compound as an amber oil (0.1g, 89%).

- 5 Mass spectrum (API^+): Found 242 (MH^+). $\text{C}_{15}\text{H}_{19}\text{N}_3$ requires 241.

^1H NMR (CDCl_3) δ : 0.30 - 0.50 (2H, m), 0.70 - 0.90 (2H, m), 1.45 (2H, br s), 2.40 - 3.00 (8H, m), 3.68 (2H, s), 7.17 (1H, d, $J = 8$ Hz), 7.32 (1H, s), 7.37 (1H, d, $J = 8$ Hz).

- 10 The following compound was prepared in a similar manner to Description 5.

(a) *trans*-2-(1-(4-Aminomethyl)cyclohexylmethyl)-7-cyano-1,2,3,4-tetrahydroisoquinoline

- 15 Mass spectrum (API^+): Found 284 (MH^+). $\text{C}_{18}\text{H}_{25}\text{N}_3$ requires 283.

Description 6

6-Cyano-1,2,3,4-tetrahydroisoquinoline

20

Prepared in a similar manner to that described in H.G. Selnick *et al.*, Synthetic Communications 25 (20) 3255 (1995).

Mass spectrum (API^+): Found 159 (MH^+). $\text{C}_{10}\text{H}_{10}\text{N}_2$ requires 158.

25

Description 7

4-(2-(7-Cyano-1,2,3,4-tetrahydro)isoquinolinyl)cyclohexanone

- 30 A mixture of 7-cyano-1,2,3,4-tetrahydroisoquinoline (2.37g, 15 mmol), 1,4-dioxaspiro-[4.5]decan-8-one (2.34g, 15 mmol) and sodium triacetoxyborohydride (4.73g, 22.5 mmol) in dichloroethane (50ml) was stirred at 20°C for 18h. Mixture was partitioned between saturated aqueous NaHCO_3 (250ml) and dichloromethane (3 x 50ml) and the combined organic extracts dried (Na_2SO_4) and evaporated *in vacuo* to give an oil. Chromatography
35 on silica with 25 - 100% ethyl acetate - hexane gradient elution gave a solid (3.53g). The latter was dissolved in water containing concentrated H_2SO_4 (1.35g, 13.5 mmol) and heated at 65°C for 18h. Mixture was cooled, then partitioned between saturated aqueous NaHCO_3 (300ml) and dichloromethane (3 x 100ml). Combined organic extracts were

dried (Na_2SO_4) and evaporated *in vacuo* to give the title compound (3.14g, 82%) as an oil.

Mass spectrum (API^+): Found 255 (MH^+). $\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}$ requires 254.

5

The following compound was prepared in a similar manner to Description 7

(a) 4-(2-(6-Cyano-1,2,3,4-tetrahydro)isoquinolyl)cyclohexanone

10 Mass spectrum (API^+): Found 255 (MH^+). $\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}$ requires 254.

Description 8

15 ***cis*- and *trans*-7-Cyano-2-(1-(4-trifluoroacetamido)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline**

A mixture of 4-(2-(6-cyano-1,2,3,4-tetrahydro)isoquinolyl)cyclohexanone 2.90g, 11.4 mmol), ammonium acetate (8.7g, 0.11 mol) and sodium triacetoxyborohydride (16.6g, 79.4 mmol) in ethanol (250ml) was heated at reflux for 3h, cooled then evaporated *in vacuo*. Residue was partitioned between saturated aqueous NaHCO_3 (300ml) and dichloromethane (3 x 100ml). Combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give an oil (2.78g). A mixture of the latter with triethylamine (2ml; 14.3 mmol) in dichloromethane (100ml) at 0°C was treated dropwise with trifluoroacetic anhydride (1.9ml, 13.5 mmol). Resulting solution was stirred at 20°C for 4h, then partitioned between saturated aqueous NaHCO_3 (300ml) and dichloromethane (3 x 100 ml). Combined organic extracts were dried (Na_2SO_4) and evaporated *in vacuo* to give an oil. Chromatography on silica with 10 - 100% ethyl acetate - hexane gradient elution gave, as the first-eluting component, the *cis*-isomer (1.58g, 38%),

30 ^1H NMR (CDCl_3) δ : 1.50 - 2.00 (8H, m), 2.48 (1H, m), 2.87 (2H, m), 2.98 (2H, m), 3.78 (2H, s), 4.09 (1H, m), 6.29 (1H, m), 7.22 (1H, m), 7.29 - 7.49 (2H, m),

and, as the second-eluting component, the *trans*-isomer (0.63g, 15%)

35 ^1H NMR (CDCl_3) δ : 1.22 - 1.44 (2H, m), 1.45 - 1.64 (2H, m), 2.05 (2H, m), 2.17 (2H, m), 2.55 (1H, tt, $J = 9, 2$ Hz), 2.84 (2H, m), 2.95 (2H, m), 3.78 (2H, s), 3.80 (1H, m), 6.15 (1H, m), 7.19 (1H, d, $J = 8$ Hz), 7.32 (1H, d, $J = 1$ Hz), 7.40 (1H, dd, $J = 8, 1$ Hz).

The following compounds were prepared in a similar manner to Description 8.

(a) *cis*-6-Cyano-2-(1-(4-trifluoroacetamido)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline

5

¹H NMR (CDCl₃) δ: 1.65 - 1.95 (8H, m), 2.47 (1H, m), 2.83 (2H, m), 1.92 (2H, m), 3.77 (2H, s), 4.05 (1H, m), 6.28 (1H, br s), 7.13 (1H, d, J = 8 Hz), 7.39 (2H, m).

(b) *trans*-6-Cyano-2-(1-(4-trifluoroacetamido)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline

10

¹H NMR (CDCl₃) δ: 1.24 - 1.62 (4H, m), 2.03 (2H, m), 2.15 (2H, m), 2.53 (1H, tt, J = 9, 2 Hz), 2.82 (2H, m), 1.86 (2H, m), 3.76 (1H, m), 3.80 (2H, s), 6.12 (1H, m), 7.09 (1H, d, J = 8 Hz), 7.35 (2H, m).

15

Description 9

***trans*-2-(1-(4-Amino)cyclohexyl)-7-cyano-1,2,3,4-tetrahydroisoquinoline**

20 A mixture of *trans*-7-cyano-2-(1-(4-trifluoroacetamido)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline (0.68g, 1.9 mmol), methanol (30ml), water (3.5ml) and anhydrous potassium carbonate (1.3g, 9.6 mmol) was heated at reflux for 3h, cooled then evaporated *in vacuo*. Residue was partitioned between saturated aqueous K₂CO₃ (50ml) and dichloromethane (3 x 50 ml), and the combined extracts were dried (Na₂SO₄) and
25 evaporated *in vacuo* to give the title compound (0.48g, 96%) as an oil.

Mass spectrum (API⁺): Found 256 (MH⁺). C₁₆H₂₁N₃ requires 255.

The following compounds were prepared in a similar manner to Description 9.

30

(a) *trans*-2-(1-(4-Amino)cyclohexyl)-6-cyano-1,2,3,4-tetrahydroisoquinoline

Mass spectrum (API⁺): Found 256 (MH⁺). C₁₆H₂₁N₃ requires 255.

(b) *trans*-2-(1-(4-(2-Amino)ethyl)cyclohexyl)-7-cyano-1,2,3,4-tetrahydroisoquinoline

35

Mass spectrum (API⁺): Found 284 (MH⁺). C₁₈H₂₅N₃ requires 283.

Description 10**4-(2-Trifluoroacetamidoethyl)cyclohexanone**

- 5 To a mixture of 8-(2-hydroxyethyl)-1,4-dioxaspiro[4.5]decane (15.5g, 83 mmol) and triethylamine (15.2ml; 0.108 mol) in dichloromethane (300ml) under argon at 0°C was added dropwise a solution of methylsulfonyl chloride (7.4ml; 96 mmol) in dichloromethane (10ml). Resulting solution was stirred at 20°C for 2h, then partitioned between saturated aqueous NaHCO₃ (500ml) and dichloromethane (3 x 50ml). Combined organic extracts
10 were dried (Na₂SO₄) and evaporated *in vacuo* to give an oil (21.8g). The latter was dissolved in toluene (50ml) and added to a solution of trifluoroacetamide anion prepared by portionwise addition of trifluoroacetamide (7.91g, 70 mmol) to a stirred suspension of sodium hydride (60%; 2.6g, 65 mmol) in dimethylformamide (50ml). The resulting mixture was stirred at 20°C for 18h, then evaporated *in vacuo*. Residue was partitioned
15 between ether (500ml) and water (350ml). Organic phase was washed with water (2 x 200ml), dried (Na₂SO₄) and evaporated *in vacuo* to give an oil (15g). Chromatography on silica with 10 - 100% ethyl acetate - hexane gradient elution gave an oil (4.96g). A solution of the latter in tetrahydrofuran (200ml) was treated with water (400ml) and concentrated H₂SO₄ (50 drops), then heated at reflux for 3h. The mixture was cooled,
20 concentrated *in vacuo* to 200ml, then extracted with dichloromethane (3 x 200ml). Combined extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give the title compound (3.72g, 19%) as a colourless solid.

25 Mass spectrum (API): Found 236 (M-H)⁺. C₁₀H₁₄F₃NO₂ requires 237.

25

Description 11***cis*- and *trans*-7-Cyano-2-(1-(4-(2-trifluoroacetamido)ethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline**

30

- A mixture of 7-cyano-1,2,3,4-tetrahydroisoquinoline (1.5g, 9.5 mmol), 4-(2-trifluoroacetamidoethyl)cyclohexane (2.25g, 9.5 mmol) and sodium triacetoxyborohydride (3.0g, 14.3 mmol) in dichloromethane (100ml) was treated with glacial acetic acid (10 drops) and stirred at 20°C for 18h. Mixture was partitioned between saturated aqueous
35 NaHCO₃ (300ml) and dichloromethane (4 x 50ml), and the combined extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give an oil (4.0g). Chromatography on silica with 10 - 100% ethyl acetate - hexane gradient elution gave, as the first-eluting component, the *cis*-isomer (1.98g, 55%)

- 40 ¹H NMR (CDCl₃) δ: 1.44 - 1.85 (11H, m), 2.45 (1H, m), 2.81 (2H, m), 2.92 (2H, m), 3.40 (2H, m), 3.72 (2H, s), 6.31 (1H, br s), 7.20 (1H, d, J = 8 Hz), 7.35 (1H, d, J = 1 Hz), 7.40 (1H, dd, J = 8, 1 Hz),

and, as the second-eluting component, the *trans*-isomer (0.92g, 26%).

¹H NMR (CDCl₃) δ: 0.95 - 1.17 (2H, m), 1.20 - 1.68 (5H, m), 1.84 - 2.07 (4H, m), 2.50 (1H, tt, J = 9, 2 Hz), 2.85 (2H, m), 2.93 (2H, m), 3.42 (2H, q, J = 7 Hz), 3.78 (2H, s),
5 6.32 (1H, br s), 7.19 (1H, d, J = 8 Hz), 7.33 (1H, d, J = 1 Hz), 7.40 (1H, d, J = 8, 1 Hz).

Description 12

trans-4-(*t*-Butyloxycarbonyl)aminomethylcyclohexanecarboxaldehyde

10

A mixture of *trans*-4-aminomethylcyclohexanecarboxylic acid (20g, 0.127 mol), methanol (250ml) and concentrated sulfuric acid (7.5ml; 0.14 mmol) was heated at reflux for 5h then evaporated *in vacuo* to give a solid. The latter was mixed with dichloromethane (250ml), triethylamine (64.5ml, 0.463 mol) and di-*t*-butyl dicarbonate (34g, 0.155 mol),
15 and the resulting solution stirred at 20°C for 18h. Mixture was partitioned between saturated aqueous NaHCO₃ (1L) and dichloromethane (3 x 200ml), and the combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give a solid (36.6g). The latter was dissolved in toluene (500ml) and cooled to -78°C under argon. A solution of diisobutylaluminium hydride in toluene (1M; 270ml) was added dropwise over 0.75h,
20 and stirring at -78°C was continued for 1h. Methanol (54.5ml) was added dropwise over 0.5h and mixture stirred at -70°C for 0.25h. The resulting solution was then poured into saturated aqueous potassium sodium tartrate (1L), and the mixture stirred vigorously for 3h. The resultant was extracted with ether (3 x 200ml) and the combined organic extracts were dried (Na₂SO₄) and evaporated *in vacuo* to give an oil (35.5g). Chromatography on
25 silica with 10 - 100% ethyl acetate - hexane gradient elution gave the title compound (20.9g, 64%) as an oil.

30

¹H NMR (CDCl₃) δ: 0.92 - 1.09 (2H, m), 1.18 - 1.50 (3H, m), 1.46 (9H, s), 1.89 (2H, m), 2.04 (2H, m), 2.19 (1H, m), 3.00 (2H, t, J = 7 Hz), 4.60 (1H, br s), 9.61 (1H, s).

Example 1

(±)-*trans*-1-((*E*)-3-(5-Indolyl)propenamido)methyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)methylcyclopropane

35

A mixture of (±)-*trans*-1-aminomethyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)methylcyclopropane (0.1g, 0.4 mmol), (*E*)-3-(5-indolyl)propenoic acid (0.09g, 0.5 mmol) 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.13g) and 1-hydroxybenzotriazole (0.06g) in dimethylformamide (1ml) and
40 dichloromethane (7ml) was shaken for 20h, then washed with water (7ml). Chromatography of the organic phase on silica, eluting with ethyl acetate in hexane 20% - 100%, gave the title compound as a colourless solid (0.11g, 66%).

Mass spectrum (API⁺): Found 411 (MH⁺). C₂₆H₂₆N₄O requires 410.

¹H NMR (CDCl₃) δ: 0.44 - 0.64 (2H, m), 0.90 - 1.00 (2H, m), 2.35 - 2.55 (2H, m), 2.75 - 3.00 (4H, m), 3.20 - 3.50 (2H, m), 3.69 (2H, s), 2.75 - 2.80 (1H, m), 6.35 (1H, d, J = 15 Hz), 6.57 (1H, m), 7.10 - 7.40 (6H, m), 7.70 - 7.85 (2H, m), 8.40 (1H, br s).

The following compounds were prepared in a similar manner to Example 1

10 (a) *trans-(E)-6-Cyano-2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline*

Mass spectrum (API⁺): Found 404 (MH⁺). C₂₅H₂₆FN₃O requires 403.

15 NMR (CDCl₃) δ: 1.25 (2H, m), 1.42 - 1.64 (2H, m), 2.00 (2H, m), 2.19 (2H, m), 2.54 (1H, m), 2.85 (4H, m), 3.81 (2H, s), 3.90 (1H, m), 4.47 (1H, d, J = 8 Hz), 6.30 (1H, d, J = 16 Hz), 7.09 (3H, m), 7.34 - 7.55 (4H, m), 7.60 (1H, d, J = 16 Hz).

20 (b) *trans-(E)-7-Cyano-2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline*

Mass spectrum (API⁺): Found 386 (MH⁺). C₂₅H₂₇N₃O requires 385.

25 ¹H NMR (CDCl₃) δ: 1.25 (2H, m), 1.44 - 1.66 (2H, m), 2.00 (2H, m), 2.16 (2H, m), 2.44 (1H, m), 2.81 (2H, m), 2.93 (2H, m), 3.76 (2H, m), 3.90 (1H, m), 5.45 (1H, d, J = 8 Hz), 6.35 (1H, d, J = 16 Hz), 7.19 (1H, d, J = 8 Hz), 7.34 (5H, m), 7.49 (2H, m), 7.62 (1H, d, J = 16 Hz).

30 (c) *trans-7-Cyano-2-(1-(4-(2-(2-indolyl)carboxamido)ethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline*

Mass spectrum (API⁺): Found 427 (MH⁺). C₂₇H₃₀N₄O requires 426.

35 ¹H NMR (CDCl₃ + CD₃OD) δ: 0.95 - 1.18 (2H, m), 1.25 - 1.48 (3H, m), 1.58 (2H, q, J = 7 Hz), 1.96 (4H, m), 2.50 (1H, m), 2.85 (2H, m), 2.94 (2H, m), 3.50 (2H, m), 3.79 (2H, s), 6.59 (1H, m), 6.89 (1H, s), 7.09 - 7.24 (2H, m), 7.29 (2H, m), 7.37 - 7.50 (2H, m), 7.65 (1H, d, J = 8 Hz), 9.84 (1H, br s).

40 (d) *trans-(E)-7-Cyano-2-(1-(4-(3-phenylpropenoyl)aminomethyl)cyclohexylmethyl)-1,2,3,4-tetrahydroisoquinoline*

Mass spectrum (API⁺): Found 414 (MH⁺). C₂₇H₃₁N₃O requires 413.

¹H NMR (CDCl₃) δ: 0.78 - 1.15 (4H, m), 1.56 (2H, m), 1.86 (4H, m), 2.31 (2H, d, J = 7 Hz), 2.69 (2H, t, J = 6 Hz), 2.93 (2H, t, J = 6 Hz), 3.27 (2H, t, J = 7 Hz), 3.59 (2H, s), 5.74 (1H, m), 6.40 (1H, d, J = 16 Hz), 7.19 (1H, d, J = 8 Hz), 7.33 (1H, s), 7.38 (4H, m), 7.51 (2H, m), 7.64 (1H, d, J = 16 Hz).

5

(e) *trans*-7-Cyano-2-(1-(4-(2-indolyl)carboxamidomethyl)cyclohexylmethyl)-1,2,3,4-tetrahydroisoquinoline

Mass spectrum (API⁺): Found 427 (MH⁺). C₂₇H₃₀N₄O requires 426.


10

¹H NMR (CDCl₃ + CD₃OD) δ: 0.85 - 1.17 (4H, m), 1.60 (2H, m), 1.90 (4H, m), 2.34 (2H, d, J = 7 Hz), 2.71 (2H, t, J = 6 Hz), 2.95 (2H, t, J = 6 Hz), 3.32 (2H, t, J = 7 Hz), 3.60 (2H, s), 6.75 (1H, m), 6.91 (1H, s), 7.07 - 7.36 (4H, m), 7.42 (2H, m), 7.64 (1H, d, J = 8 Hz), 9.95 (1H, br s).

15

Claims :

1. A compound of formula (I):



Formula (I)

5 wherein:

R¹ represents a substituent selected from: a hydrogen or halogen atom; a hydroxy, cyano, nitro, trifluoromethyl, trifluoromethoxy, trifluoromethanesulfonyloxy, pentafluoroethyl, C₁₋₄alkyl, C₁₋₄alkoxy, arylC₁₋₄alkoxy, C₁₋₄alkylthio, C₁₋₄alkoxyC₁₋₄alkyl, C₃₋₆cycloalkylC₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkoxycarbonyl, C₁₋₄alkylsulphonyl, C₁₋₄alkylsulphonyloxy, C₁₋₄alkylsulphonylC₁₋₄alkyl, arylsulphonyl, arylsulphonyloxy, arylsulphonylC₁₋₄alkyl, C₁₋₄alkylsulphonamido, C₁₋₄alkylamido, C₁₋₄alkylsulphonamidoC₁₋₄alkyl, C₁₋₄alkylamidoC₁₋₄alkyl, arylsulphonamido, arylcarboxamido, arylsulphonamidoC₁₋₄alkyl, arylcarboxamidoC₁₋₄alkyl, aroyl, aroylC₁₋₄alkyl, or arylC₁₋₄alkanoyl group; a group R³OCO(CH₂)_p, R³CON(R⁴)(CH₂)_p, R³R⁴NCO(CH₂)_p or R³R⁴NSO₂(CH₂)_p where each of R³ and R⁴ independently represents a hydrogen atom or a C₁₋₄alkyl group or R³R⁴ forms part of a C₃₋₆azacycloalkane or C₃₋₆(2-oxo)azacycloalkane ring and p represents zero or an integer from 1 to 4; or a group Ar³-Z, wherein Ar³ represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring and Z represents a bond, O, S, or CH₂;

s represents an integer from zero to 2 and r represents an integer from 1 to 4, such that the sum of s + r is 1 to 4;

t represents an integer from zero to 1 and u represents an integer from zero to 2;


R² represents a hydrogen atom or a C₁₋₄alkyl group;

q is 1 or 2;

A represents a group of the formula (a), (b) or (c):



(a)



(b)



(c)

wherein

Ar represents an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; or an optionally substituted bicyclic ring system;

Ar¹ and Ar² each independently represent an optionally substituted phenyl ring or an optionally substituted 5- or 6- membered aromatic heterocyclic ring; and

Y represents a bond, -NHCO-, -CONH-, -CH₂-, or -(CH₂)_mY¹(CH₂)_n-, wherein Y¹ represents O, S, SO₂, or CO and m and n each represent zero or 1 such that the sum

of $m+n$ is zero or 1; providing that when A represents a group of formula (a), any substituent present in Ar ortho to the carboxamide moiety is necessarily a hydrogen or methoxy group;

and salts thereof.

5


2. A compound according to claim 1 wherein q represents 1.

3. A compound of formula (I) which is:

- 10 (\pm) -*trans*-1-((*E*)-3-(5-Indolyl)propenamido)methyl-2-(2-(7-cyano-1,2,3,4-tetrahydro)isoquinolyl)methylcyclopropane
trans-(*E*)-6-Cyano-2-(1-(4-(3-(4-fluoro)phenylpropenoyl)amino)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline
trans-(*E*)-7-Cyano-2-(1-(4-(3-phenylpropenoyl)amino)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline
 15 *trans*-7-Cyano-2-(1-(4-(2-(2-indolyl)carboxamido)ethyl)cyclohexyl)-1,2,3,4-tetrahydroisoquinoline
trans-(*E*)-7-Cyano-2-(1-(4-(3-phenylpropenoyl)aminomethyl)cyclohexylmethyl)-1,2,3,4-tetrahydroisoquinoline
 20 *trans*-7-Cyano-2-(1-(4-(2-indolyl)carboxamidomethyl)cyclohexylmethyl)-1,2,3,4-tetrahydroisoquinoline
 or a salt thereof.

4. A process for preparing compounds of formula (I) which process comprises

25 (a) reacting a compound of formula (V):



Formula (V)

30

with a compound of formula (VI):


A-COX

35

Formula (VI)

wherein A is as hereinbefore defined and X is a halogen atom or the residue of an activated ester;

40 (b) to prepare a compound of formula (I) wherein R^1 is Ar^3-Z and Z is a bond, reacting a compound of formula (VII):




Formula (VII)

5

wherein one R^{1a} represents a group W wherein W is a halogen atom or a trifluoromethylsulphonyloxy group, or W is a group M selected from a boron derivative or a metal function, and when q is 2 the other R^{1a} is R^1 ; with a compound Ar^3-W^1 , wherein W^1 is a halogen atom or a trifluoromethylsulphonyloxy group when W is a group M or W^1 is a group M when W is a halogen atom or a trifluoromethylsulphonyloxy group;

10

(c) to prepare a compound of formula (I) wherein R^1 is Ar^3-Z and Z is O or S, reacting a compound of formula (VIII):




15

Formula (VIII)

wherein one R^{1b} represents a group ZH and when q is 2 the other R^{1b} represents R^1 ; with a reagent serving to introduce the group Ar^3 ;

20

(d) to prepare a compound of formula (I) where Y is a bond, reaction of a compound of formula (IX):



Formula (IX)

25

wherein R^1 , R^2 , Ar^1 and W are as hereinbefore defined, with a compound Ar^2-W^1 , wherein W^1 is a halogen atom or a trifluoromethylsulphonyloxy group when W is a group M, or W^1 is a group M when W is a halogen atom or a trifluoromethylsulphonyloxy group.

(e) interconversion of one compound of formula (I) to a different compound of formula (I);

30

(f) where appropriate, separation of enantiomers, diastereoisomers, or *cis*- and *trans*- isomers of compounds of formula (I), or intermediates thereto, by conventional methods;

and optionally thereafter forming a salt of formula (I).

5 5. A pharmaceutical composition comprising a compound of formula (I) as claimed in any of claims 1 to 3 or a physiologically acceptable salt thereof and a physiologically acceptable carrier therefor.

5 6. The use of a compound of formula (I) as claimed in any of claims 1 to 3 or a physiologically acceptable salt thereof in the manufacture of a medicament for the treatment of a condition which requires modulation of a dopamine receptor.

10 7. Use according to claim 6 wherein the dopamine receptor is a dopamine D₃ receptor.

 8. Use according to claim 6 or claim 7 wherein a dopamine antagonist is required.

15 9. Use according to any of claims 6 to 8 wherein the condition is a psychotic condition.

20 10. A method of treating a condition which requires modulation of a dopamine receptor which comprises administering to a subject in need thereof an effective amount of a compound of formula (I) as claimed in claim 1 or a physiologically acceptable salt thereof.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 98/02584

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07D217/04 C07D401/12 A61K31/47		
According to International Patent Classification(IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07D A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 02246 A (BASF AG) 1 February 1996 see the whole document ---	1,5-10
A	US 5 294 621 A (RUSSELL RONALD K) 15 March 1994 cited in the application see the whole document ---	1,5-10
P,A	WO 98 06699 A (SMITHKLINE BEECHAM PLC) 19 February 1998 see claims ---	1,5-10
P,A	WO 97 43262 A (SMITHKLINE BEECHAM PLC) 20 November 1997 see claims -----	1,5-10
<div style="display: flex; justify-content: space-between;"> <input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex. </div>		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold;">20 August 1998</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold;">27/08/1998</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold;">Henry, J</div>

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 98/ 02584

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 10
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim 10
is directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 98/02584

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9602246 A	01-02-1996	DE 4425146 A AU 3111495 A CA 2195242 A CN 1152870 A CZ 9700096 A EP 0771197 A FI 970148 A HU 77608 A JP 10502658 T NO 970163 A SI 9520084 A	18-01-1996 16-02-1996 01-02-1996 25-06-1997 13-08-1997 07-05-1997 14-01-1997 29-06-1998 10-03-1998 14-03-1997 31-08-1997
US 5294621 A	15-03-1994	NONE	
WO 9806699 A	19-02-1998	AU 4204697 A	06-03-1998
WO 9743262 A	20-11-1997	AU 2897497 A	05-12-1997